



Educating Health Workers about Dealing with Seasonal Infections

1Moshawi Ali Gosairy, 2Akram Mohammed Ali Sahloli, 3Zahra Mohammed Eisi Kuayti, 4Areej Shoui Ali Bahari, 5Mohammed Abdullah Ahmad Alzhrani, 6Barakat Khudaysh Barakat Alzahrani, 7Abdulaziz Ali Yasin Alkhayri, 8Haya Mubarak Masoud Al Mohammed, 9Afaf Taher Nafe, 10Haifa Falah Alotaibe

1Pharmacy Technician, Edabi Hospital

2Nurse Technician, Edabi Hospital

3Nurse Technician, Khadera Primary Health Care

4Nursing Specialist, Sabya General Hospital

5Nursing Technician, Al-Mudhaylif Primary Health Care Center

6Nursing Technician, Al-Mudhaylif Primary Health Care Center

7Nursing Technician, Al-Mudhaylif Primary Health Care Center

8Health Assistant, Irada Complex For Mental Health And Addiction

9Health Assistant, Irada Complex For Mental Health And Addiction

10Nursing Technician, Diriyah Hospital

1. Introduction to Seasonal Infections

In communities across the globe, healthcare workers become increasingly vigilant as summer transitions to fall. For it is during this period that various seasonal infections, notable among them the flu, circulate and strike down individuals, resulting in significant morbidity, bed loss, outpatient and emergent visits, and in some instances horrific complications and loss of life. Healthcare workers such as doctors, nurses, and allied care professionals are reminded in various ways to communicate the importance of vaccination and hygiene to avoid these entirely preventable problems. Nonetheless, when we reflect on the education and training of this cadre of professionals, it is surprising to see the relative absence of specific curricular content directed towards the comprehensive understanding of seasonal respiratory infections. While it is true that areas of content pertaining to microbiology, infectious diseases, and public health in general do appear in some guise in medical, nursing, and other allied health curriculums, these often do not specifically encompass practical knowledge that healthcare workers can use on an everyday basis to diagnose, manage, and curtail community transmission of these infections.



Methods

Standardized educational programs are effective at improving health workers' knowledge about influenza. Training initiatives in response to outbreaks are as effective as longer programs. However, longer training periods may reduce response time and participation. Teaching health and hygiene alongside each other is necessary for effectiveness. Providing relevant information and problem-solving resources improves training outcomes. Personal hygiene can be addressed at work or at home.

Conclusion

It is remembered how sharply it was realized during the discussion of the report on live education about influenza that educational events carried out over the past three years were aimed only at increasing the motivation for vaccination and the proportion of vaccinated. There was little talk about the basic issue-educating health workers in facets of prevention, clinical reasoning and management under existing medical guidelines. This area really is and is very important. Everyone called him symbolic. Although with the full understanding of the capabilities of information and communication technologies and the arrested activities needed by health workers, it was recognized that the issue would remain relevant for a very long time. It is very important and timely that experts have turned to this problem and developed a strategy that will undoubtedly allow countries to effectively cultivate health workers in facets of seasonal prevention through the use of each of the points of the strategy presented.

Introduction

One of the basic tenets of dealing with a pandemic is to have people who can provide education in relevant aspects of respiratory physiology, infection refractoriness, conditions that exacerbate the severity of respiratory infections, and hygienic measures that can reduce the likelihood of respiratory microbial exposure. Given the crucial importance of this borderland between the science of pandemics and human communication, it is most appropriate to have health workers who are knowledgeable in these aspects that play such a crucial role in the spread of infections. Educating workers involved in infection control is not a simple matter; while the curriculum may lower barriers such as the unfamiliarity of students with aspects dealing with the science underlying so-called common sense rules for avoiding infections they may have heard for much of their lives, it is unrealistic to believe that a relative ocean of pre-existing ideas can be replaced by a newer set based on widely held laboratory research results. This is particularly true for those who provide direct patient care at the coalface of the infection battle. The paradox is one where, quite correctly, health workers provide support to the inculcation of values that the science, while being much less frequently taught, indicates are wise approaches. The lessons in etiology that are often transmitted in science textbooks can also prepare students to point out the limitations of commercially available health materials that either cannot work or are so poorly made they



do not work. Those who develop educational materials need to be able to anticipate questions and statements that their introductory segments may reasonably prompt.

1.1. Definition and Types of Seasonal Infections

While we all know that infectious diseases exist in all parts of the world, knowledge about the characteristics of organisms and environmental conditions can reduce or avoid infection. Approximately 22% of the total disease burden in relation to death, disability, and loss of working days in India has been estimated to be due to preventable infectious diseases. India has reported a 51.1% mortality incidence due to infectious and parasitic diseases. Prevention is always better than treatment. The knowledge and understanding of the impact of environmental factors are necessary to aid in the prevention of disease. There have been enormous strides in the area of prevention in the modern world with the development of modern science and technology. Specific knowledge of the season and identifying the phenomenon between season and communicable diseases is important for the prevention of season-specific infections.

Microorganisms are generally present in the environment. Their growth, distribution, and pathogenicity depend on specific environmental factors. The environmental hazards in health and the inherent relationship between human habitat can be linked to seasonal changes. These environmental factors are temperature, radiation, humidity, salinity, and other related climatic conditions. The biotic and abiotic components are present in the environment in conjunction. Humans are homeothermic organisms, while the growth and activity of microorganisms are influenced by environmental temperature. Cloths, towels, air, water, and soil are vital media for the transmission of pathogenic organisms. Due to the human habitat and their ingestion, pathogenic organisms enter the oral cavity. Disease depends not only on specific pathogenic organisms and immune status but also on the above intrinsic interactions in the environment. It is established that indigenous dwellers exhibit decreased susceptibility to infectious diseases prevalent in the region. Seasonal variation in diseases, including seasonal outbreaks, was recorded. These aspects related to season-specific diseases and seasonal outbreaks are discussed in the following section.

1.2. Impact on Public Health

For more than 50 years, influenza has been a notifiable disease in Germany. In spite of the labor-intensive detection process required, the documented rates are still extant, making influenza the most important epidemic infection besides gastroenteritis and bronchitis during the colder months. Social costs for society due to influenza are by far higher. Influenza is not only one of the most heavily contagious viral diseases, but is also statistically more dangerous than many other agents. With its typical secondary bacterial infections, in particular pneumonia, it really has the potential to depopulate the susceptible risk groups. Obesity is on the increase as a risk factor with an alarming exponential tendency, particularly



in younger persons. Pregnant women and their neonates are especially severely affected, both in absolute numbers and relative findings during the pandemic seasonal peak. Also, persons over age 65 and patients with pulmonary diseases have to be classified as risk groups. Currently, vaccine recommendations are pronounced for these patients, as well as for health care personnel directly in contact with the patients of the risk groups. (Ullrich et al.2021)(Oh et al.2021)(van et al.2023)

New educational strategies are required in order to interest doctors and nurses in these educational offers. Only experienced nursing staff has access to some necessary handcrafted tricks, such as sponge baths, cough assistance, and adapted mobilization programs. This know-how has to be handed down from experienced to novice staff at a low threshold level. Also, laypersons could profit from educational packages concerning this seasonal infection. In particular, the school medicine departments would be more than willing to offer training courses for their students in health care facilities and in universities. Educational medical games, problem-based learning, or clinical education based on serious events provide more plausibility and higher learning success in a more engaging way. Also, for simple sanitary behavior such as proper hand washing, adapted sneezing, and coughing, entertaining forms of educational offers give a better incentive and learning success.

2. Role of Health Workers in Seasonal Infection Management

Despite advanced techniques in medicine, the training of proper health workers is neglected. Both the Head Office and Peripheral Health Workers should know the management aspects of seasonal infections. HQ personnel should be informed, trained, and reminded to focus on seasonal infection relief in a timely manner and strongly suggested to support health workers in seasonal infection management, as routine health care work would run over the usual speed during periodic peak seasonal infection periods. Allowing extra staff support, communication, transportation, and basic logistics would be helpful to run the services effectively and achieve better results.

Earlier assisted management and relief would prevent suffering, which would provide significant relief to families and the local economy. All such activities boost the social respect of health care providers. This is essential while the Health and Family Welfare Service is under serious pressure due to various negative reports. Health care workers should lead with love and respect, not with enforcement. Only then will meaningful change occur in public cooperation, and trust-building will proceed at the proper pace.

2.1. Key Responsibilities and Duties

The goal of educating health workers about dealing with seasonal infections is to prepare them to handle all levels of disease prevention and treatment. This system includes instructions to help the health workers recognize typical signs of seasonal infections, as well as suggestions on how to work together with community volunteers to decrease health



threats. The program objective is to ensure that health workers serving in rural areas are well-prepared to control and handle common health problems found in their community once the more senior health staff are not immediately available to give help and advice. This is a vital task. In a year, a number of children will be in danger of dying or becoming very sick from seasonal infections. To protect the life and health of the rural community from these known dangers, the public must be served by people who are trained and capable of handling everyday problems. These people are the general health care workers such as the health guide, family planning assistant, and traditional practitioners who generally reside in the community and are known to individuals, families, and community organizations. These people have an additional and important job to do, a job that cannot be postponed, so by teaching them and their community workers how to do it effectively, all members of the rural community will gain greater confidence in their abilities concerning treatment and prevention of seasonal infections.

3. Educational Strategies for Health Workers

In many countries, health personnel – practitioners, nurses, other care providers, and public health staff face 'influenza seasons,' which are peaks of respiratory and influenza-like illness (ILI) cases. Daunting work pressures on these health workers may be exacerbated by serious illness and staffing deficiencies related to illness, which would be further complicated in the case of a pandemic situation. While 'strains' of influenza viruses must always be monitored for pandemic potential, newly appearing viruses show potential for pandemic spread; the influenza these strains cause may be clinically variable. (Hudu et al.2024)(Ehrlich et al., 2020)(Jacek and Mastalerz-Migas2022)(Rios et al.2020)

In our studies, sustained infection of personnel in health settings with 'seasonal' viruses was suggested to help produce novel variants of human-targeted viruses, and allowing these genetically variable viruses to co-circulate with seasonal H1N1 and H3N2 viruses was further suggested to help make novel actions of the pandemic virus less virulent. Hence, it would be desirable to manage the effects of mild influenza infections in health care settings to reduce work accumulation and individual illness. Educational strategies to reduce absenteeism and 'presenteeism' during influenza seasons and to control apparent dissemination of virus varieties are discussed.

3.1. Training Programs and Workshops

A variety of training programs and workshops take place in countries in preparation for seasonal infections. These training programs include basic information about transmission, vaccination, and other prevention and treatment methods. Traditional winter disease accessories that are distributed, such as scarves, are often included. The programs take place at different times, depending on the roles of the workers: early-bird teachers, neighborhood mothers or community care educators, and health workers, such as pediatric and family



physicians, pediatric nurses, and support staff of family health centers. At the second level of health care, pediatric pulmonologists are also involved. In countries where health workers are employed by the private sector, influenza vaccination campaigns specifically target general practitioners and pediatricians.

What is unique about the programs in Turkey is that the majority of teachers are women, and these women frequently live with extended families. Because of these roles and responsibilities, teachers interact with parents to organize holiday activities for children or with mothers to give advice about health-related issues. A program in early childhood education offers the opportunity to provide education about upper respiratory tract infections with the assistance of a teaching program about common viral, bacterial, and chlamydia infections. It also offers the opportunity to educate teachers about basic information such as hand hygiene, appropriate methods of greeting one another during the flu season and after it has ended, and prevention and treatment methods. Programs for teachers are offered free of charge as part of the winter disease program.

Educating health workers about seasonal infections—especially those of the upper respiratory tract—should be seen as important in the struggle to combat community infections, especially those that are viral in nature, from primary health care to the tertiary level.

4. Preventive Measures and Best Practices

Best knowledge practices in infectious disease risk and prevention will offer a comprehensive orientation about modes of illness prevention among vulnerable populations, which include children, elderly people, low-income, and ethnic communities in preventing exposure to the germs that cause infections. Influenza vaccination is safe and has been associated with a reduced burden of respiratory infections, reduced antibiotic use, fewer missed days at work, and lower hospitalization rates. Preventive measures include: 1. routine handwashing with soap and tap water, especially before eating and food preparation; 2. routine handwashing with soap and tap water, especially after use of the toilet, changing diapers, and handling garbage; 3. cough etiquette such as covering the mouth and nose during a cough using a tissue or the upper portion of the sleeve, elbow, or arm if no tissue is available; 4. appropriate immunization for epidemic infections. Bloodborne infection prevention and post-exposure protocols relevant to disease-preventive behaviors are essential to learn about for added protection when working as a health care worker. General principles: The purpose of a respiratory hygiene and cough etiquette program is to prevent the spread of infectious respiratory agents in institutional settings.

4.1. Vaccination and Immunization

Seasonal flu is a highly infectious, easily transmitted, vaccine-preventable respiratory disease caused by influenza viruses. Almost all countries have routine seasonal influenza vaccination policies for high-risk groups and for health professionals. The general objectives of these



recommendations are the prevention of nosocomial infection in health care. Protecting healthcare professionals from many illnesses ensures that they can maintain staffing levels, functionality, and protection for themselves and their families, while avoiding the consumption of health resources. In all cases, the performance of quantitative serological tests is recommended to ensure the achievement of an adequate immune response to the vaccine.

Health professionals are informed about the severity of the consequences of not vaccinating themselves and the population that has contact, enabling them to make the best decisions. Income is often a critical factor associated with the non-adherence of workers to vaccination signals. Inadequate perceptions are always associated with the protection against the vaccine. If the professionals were aware and well-informed, they would not have been at risk. The uninformed believe that they only have contact with healthy people and should only be vaccinated when they fall within the groups mentioned above.

5. Conclusion and Future Directions

In conclusion, health education is an important preventive measure for dealing with seasonal infections such as influenza. If health workers use this knowledge to modify certain risky behaviors that facilitate the spread of the infection, such as washing their hands adequately and keeping their distance, then the spread of the infection could be significantly minimized. Future directions could include prospective investigations in various health areas and at different levels of preparation for the prevention of seasonal infections, as well as a system of continuous review of the consequences of these educational interventions, such as the incidence rate of sick leaves, costs, etc., as a feasible approach to emerging problems linked to potential co-circulation with new viruses or similar viruses.

References:

1. Ullrich, A., Schranz, M., Rexroth, U., Hamouda, O., Schaade, L., Diercke, M., & Boender, T. S. (2021). Impact of the COVID-19 pandemic and associated non-pharmaceutical interventions on other notifiable infectious diseases in Germany: An analysis of national surveillance data during week 1–2016–week 32–2020. *The Lancet Regional Health–Europe*, 6. [thelancet.com](https://www.thelancet.com)
2. Oh, D. Y., Buda, S., Biere, B., Reiche, J., Schlosser, F., Duwe, S., ... & Dürrwald, R. (2021). Trends in respiratory virus circulation following COVID-19-targeted nonpharmaceutical interventions in Germany, January–September 2020: Analysis of national surveillance data. *The Lancet Regional Health–Europe*, 6. [thelancet.com](https://www.thelancet.com)
3. van de Berg, S., Charles, T., Dörre, A., Katz, K., & Böhm, S. (2023). Epidemiology of common infectious diseases before and during the COVID-19 pandemic in Bavaria, Germany, 2016 to 2021: an analysis of routine surveillance data. *Eurosurveillance*, 28(41), 2300030. [eurosurveillance.org](https://www.eurosurveillance.org)



4. Hudu, S. A., Jimoh, A. O., Alqtaitat, A., & Fayig, E. (2024). The Role of Seasonal Influenza in Compounding the Outbreak of Infectious Diseases: A Critical Review. *Biomedical and Pharmacology Journal*, 17(1), 1-13. biomedpharmajournal.org
5. Ehrlich, H., Boneva, D., & Elkbuli, A. (2020). The intersection of viral illnesses: A seasonal influenza epidemic amidst the COVID-19 pandemic. *Annals of Medicine and Surgery*. sciencedirect.com
6. jacek JEdrzejek, M., & Mastalerz-Migas, A. (2022). Seasonal influenza vaccination of healthcare workers: a narrative review. *International Journal of Occupational Medicine and Environmental Health*, 35(2), 127. nih.gov
7. Rios, P., Radhakrishnan, A., Thomas, S. M., Darvesh, N., Straus, S. E., & Tricco, A. C. (2020). Guidelines for preventing respiratory illness in older adults aged 60 years and above living in long-term care: A rapid review of clinical practice guidelines. *medRxiv*, 2020-03. medrxiv.org