



Evaluating Structural Entropies and Indices in Triangular and Rhombohedral Phases of Bilayer Germanium Phosphide

Muhammad Akmal¹, Hifza Iqbal¹, Muhammad Kamran Naseer²
Muhammad Saqlain Zakir³, Muhammad Reza Farahani⁴, Mehdi
Alaeiyan⁴, Murat Cancan⁵

¹Department of Mathematics and Statistics, The University of Lahore, Defence Road Campus, Lahore, Pakistan, e-mails: akmalsial1194@gmail.com¹, Iqbalhifza3@gmail.com¹,

²Department of Mathematics, Lahore Leads University, Lahore, Pakistan. e-mail: kamrannaseer92@gmail.com²

³Department of Mathematics, COMSATS University Islamabad, Sahiwal Campus, Sahiwal, 57000, Pakistan. e-mail: saqlainsaqisial786@gmail.com³.

⁴Department of Mathematics and Computer Science, University of Science and Technology (IUST), Narmak, Tehran, 16844, Iran. e-mail: farahani@mathdep.iust.ac.ir⁴, alaeiyan@iust.ac.ir⁴

⁵Faculty of Education, Van Yuzuncu YI University, Zeve Campus, Tuba, 65080, Van, Turkey e-mail: mcancan@yyu.edu.tr⁵

Abstract— Topological indices are molecular descriptors that encode valuable information about a chemical structure. These indices enable researchers to calculate numerical values that correlate with various physical properties of chemicals, allowing for the prediction and analysis of their behaviour and characteristics. In this study, we investigate the triangular and rhombohedral bilayers of germanium phosphide, leveraging topological indices and entropy calculations. Our numerical analysis and graphical representations reveal insightful trends and patterns, providing valuable information on the validity of these indices for predicting the potential of these chemical structures for sustainable material production. This work demonstrates the power of topological indices and entropy calculations in accelerating material discovery, enabling the creation of innovative technologies with a reduced environmental footprint.

Keywords—Bilayer Germanium Phosphide; Entropy; Topological Indices; Structural Analysis; Semiconductor Applications.

1. INTRODUCTION

Within mathematical chemistry, the discipline of chemical graph theory applies graph theory to the modeling and analysis of chemical structures or networks. This method uses a graph to illustrate the structure of a chemical molecule, with atoms shown as points (nodes) and the



bonds that connect them as lines (edges). Through the use of graph theory ideas, scientists can extract many numerical parameters from these graphical representations that are referred to as topological parameters. These parameters are graph invariants, which means that they do not change based on how the structure is shown visually Ali et al., [1].

While Baig et al., [2] computed various indices for DSL (dominating set of a line graph).

In chemical graph theory, several topological characteristics are calculated to clarify physicochemical features, including configuration, enthalpy, and stability [3,4,5]. Various methods are used to derive these values; some rely on the number of edges that connect a vertex, while others measure the distance between a vertex and a particular point. In chemical graph theory, neighborhood indices are quantitative metrics that are used to describe the topological structure of molecules. They shed light on the symmetry and connectivity patterns of molecular graphs, which are essential for comprehending their characteristics and actions in a variety of applications. Here is a detailed introduction to some neighborhood indices: ABC_4 index, GA_5 index, and Sanskruti index [6,7]. Simon raj and George [10] investigated the physicochemical characteristics of dominating silicate networks. These factors have been the subject of several study articles, which have provided insightful information about various chemical structures, nanotubes, and networks, clarifying their physicochemical characteristics, including electron negativity and electron affinity.

The ABC_4 index is a neighborhood degree based index, this index has been utilized in the study of molecular descriptors and has shown applications in QSAR (Quantitative Structure-Activity Relationship) studies [8]. The GA_5 index, also known as the fifth Geometric-Arithmetic index, is another neighborhood degree based index used to quantify the molecular structure [9]. This index provides information about the symmetry and complexity of molecular graphs [11]. The Sanskruti index is a relatively newer neighborhood index in chemical graph theory. It incorporates topological features and has been applied in studies involving molecular descriptors and structure-property relationships [12].

$$N_u = \sum_{u \in \eta(v)} P_u \quad (1)$$

Where N_u represent open neighborhood degree of the vertex u.

2. PRELIMINARIES

Definition 2.1 In their 2010 paper, Ghorbani and Hosseinzadeh [17] presented the fourth iteration of the atomic-bond connectivity index (ABC_4) and the fifth iteration of the geometric arithmetic index (GA_5) for graph G as follows,

$$ABC_4 = \sum_{u_i, v_i \in E(G)} \sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}, \quad (2)$$



$$GA_5 = \sum_{u_i, v_i \in E(G)} \frac{2 \sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \quad (3)$$

Definition 2.3 In 2017, Hosamani introduced the Sanskruti index [6] for a molecular graph G , denoted by $S(G)$ and defined as,

$$S(G) = \sum_{u_i, v_i \in E(G)} \left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \quad (4)$$

Shannon first introduced the concept of entropy in his seminal work [13] in 1948. Entropy quantifies the unpredictability or uncertainty of information content within a system, often measured through the entropy of a probability distribution. Over time, this notion found applications in graph theory and chemical networks, offering insights into their structural characteristics. The utilization of graph entropies has seen a surge in various fields including biology, chemistry, ecology, and sociology. Within these domains, the degree of each atom holds significant importance. Both graph theory and network theory have extensively researched invariants, serving as functional information measures in scientific investigations. Zakir et al., [23] computed Weighted Entropy Based on Topological Indices for Triazine-Based Dendrimer

In this article, we focus on exploring graph entropy measures applied to germanium phosphide and computed, the 4th atomic bond connectivity entropy, the 5th geometric arithmetic entropy, and the Sanskruti entropy, utilizing their respective indices. Our approach draws inspiration from the concept of entropy elucidated in Manzoor et al., [14].

3. Applications of Entropy

In the context of a bilayer germanium phosphide structure, graph entropy plays a pivotal role in analyzing the structural information inherent in the system. Specifically, distance-based entropy measures are instrumental in various aspects, ranging from mathematical analyses to investigations in biology, chemical graph theory, and organic chemistry, tailored to the unique characteristics of the germanium phosphide bilayer. Drawing on Shannon's entropy concept, the structural complexity of the bilayer structure is enhanced by incorporating topological indices. These indices serve as invaluable molecular descriptors, particularly pertinent in studies concerning the properties and behaviours of the germanium phosphide bilayer, such as its electronic structure, vibrational properties, and interlayer interactions.

Shannon's seminal work in 1948 laid the groundwork for modern information theory, which found early applications in linguistics and electrical engineering before branching out into biology and chemistry. Early studies, like those in 1953, [20] paved the way for applying



information theory to address structural complexities in materials such as the bilayer germanium phosphide [21].

Further advancements in understanding the structural information content of the bilayer germanium phosphide emerged in 2004, leveraging Shannon's entropy formulas. Additionally, contributions from researchers like Rashevsky and Trucco in the 1950s [24] are pertinent to these applications, providing foundational insights that continue to inform contemporary analyses of the bilayer structure.

Graph entropy measures are widely employed in biology, computer science, and structural chemistry, and tailored applications in studying the bilayer germanium phosphide structure. These measures facilitate a comprehensive understanding of the system's properties and behaviours, aiding in tasks such as characterizing its electronic band structure, predicting its optical properties, and exploring its potential applications in nanoelectronics and optoelectronics. Overall, the applications of graph entropy in the context of a bilayer germanium phosphide structure encompass a broad spectrum of analyses, ranging from fundamental structural characterization to predictive modelling and materials design, with implications for advancing various technological frontiers.

4. Neighborhood Degree-Based Entropy

In 2015, Chen et al., [25] introduced the concept of entropy for an edge-weighted graph G denoted as $G = (V_G, E_G; (u_i, v_j))$

Here, V_G represents the set of vertices, E_G represents the set of edges, and (u_i, v_j) represents the weight associated with the edge (u_i, v_j) . The entropy of an edge-weighted graph is defined as follows, Chen et al., (2021) introduced the probability density function of a simple connected graph G given by,

$$P_{ij} = \frac{w(xy)}{\sum w(xy)} \quad (5)$$

Cao et al., (2017) introduced degree base entropy for any graph G which is denoted and defined as,

$$I(G, w) = - \sum P_{ij} \log(P_{ij}) . \quad (6)$$

Chen et al., (2014) proposed the definition of entropy of an edge-weighted graph.



$$ENT_{I(G)} = \log I(G) - \frac{1}{I(G)} \log \left\{ \prod_{u_i, v_j \in E_G} \left[\frac{N_{u_i} + N_{v_j}}{N_{u_i} \times N_{v_j}} \right]^{\left[\frac{N_{u_i} + N_{v_j}}{N_{u_i} \times N_{v_j}} \right]} \right\} \quad (7)$$

Where $I(G)$ represents the value of the topological index used to calculate the entropy of graph G . Manzoor et al., (2020) introduced the following entropies by using the above formula, The entropy of fourth atom-bond connectivity,

$$ENT_{ABC_4}(G) = \log(ABC_4(G)) - \frac{1}{ABC_4(G)} \log \left\{ \prod_{u_i, v_j \in E_G} \left[\frac{\sqrt{N_{u_i} + N_{v_j} - 2}}{N_{u_i} N_{v_j}} \right]^{\left[\sqrt{\frac{N_{u_i} + N_{v_j} - 2}{N_{u_i} N_{v_j}}} \right]} \right\} \quad (8)$$

Fifth geometry arithmetic entropy:

$$ENT_{GA_5}(G) = \log(GA_5(G)) - \frac{1}{GA_5(G)} \log \left\{ \prod_{u_i, v_j \in E_G} \left[\frac{2 \sqrt{N_{u_i} N_{v_j}}}{N_{u_i} + N_{v_j}} \right]^{\left[\frac{2 \sqrt{N_{u_i} N_{v_j}}}{N_{u_i} + N_{v_j}} \right]} \right\}, \quad (9)$$

Sanskriti entropy:

$$ENT_{S(G)} = \log(S(G)) - \frac{1}{S(G)} \log \left\{ \prod_{u_i, v_j \in E_G} \left\{ \frac{N_{u_i} \times N_{v_j}}{N_{u_i} + N_{v_j} - 2} \right\}^{\left\{ \frac{N_{u_i} \times N_{v_j}}{N_{u_i} + N_{v_j} - 2} \right\}^3} \right\} \quad (10)$$

5. Germanium Phosphide

The emergence of graphene signalled a breakthrough in the field of two-dimensional (2D) crystals, leading to a broad spectrum of materials with critical electronic characteristics required for Nanoelectronics. Semiconductors like transition metal dichalcogenides (TMDCs) and Dirac semimetals like graphene, silicene, and germanene have found useful uses in nanotechnology, especially in optoelectronics, where the first applications have been reported based on TMDCs. Every element in Group 14 of the periodic table has semi-metallic characteristics. In electronics, combining Group 14 elements with Phosphorene, an allotropic



form of Phosphorus (P), has produced positive outcomes. In 1970, a layered substance called Germanium Phosphide (GeP_3) with a stoichiometric of Phosphorene (P) and Germanene (a single-layer material composed of Germanium atoms, Ge) was initially identified. GeP_x is frequently found in three forms. Structure of $2LGeP_3$ This article uses geometrical forms such as triangles and rhombus, to illustrate the structure of GeP_3 in two distinct ways, designating as triangulate, rhombohedral, the corresponding forms of these formations. These structural forms are widely recognized in the field of nanotechnology. In the triangle, $2LGeP_3$ configuration, the bilayer structure of germanium phosphide ($2LGeP_3$) is depicted with a triangular shape. This configuration encompasses a total of $3n^2 + 9n$ edges, where n represents the number of hexagons present in the base of the monolayer $1LGeP_3$ structure, with n being greater than 2.




Figure 1: Triangular Bilayer Germanium Phosphide for $n = 4$

Neighborhood Degree-Based Partition

Now, utilizing the definition (1) of N_u , we propose a distinct edge partition for the edge set of the rhombohedral bilayer Germanium Phosphide. This partition is based on the sum degree of neighbourhood vertices associated with each edge. We have determined that for ($n > 3$), the edge partition based on neighbourhood characteristics for the edge set of ($T2LGePn3$) comprises 22 sub-partite sets, as described below.

$$\eta_1 = (u, v) | N_u = 4 \wedge N_v = 7, \quad \eta_2 = (u, v) | N_u = 10 \wedge N_v = 12$$

$$\eta_3 = (u, v) | N_u = 6 \wedge N_v = 6, \quad \eta_4 = (u, v) | N_u = 9 \wedge N_v = 13$$

$$\eta_5 = (u, v) | N_u = 5 \wedge N_v = 10, \quad \eta_6 = (u, v) | N_u = 10 \wedge N_v = 13$$

$$\eta_7 = (u, v) | N_u = 6 \wedge N_v = 10, \quad \eta_8 = (u, v) | N_u = 10 \wedge N_v = 16$$

$$\eta_9 = (u, v) | N_u = 6 \wedge N_v = 9, \quad \eta_{10} = (u, v) | N_u = 12 \wedge N_v = 16$$



$$\begin{aligned}
 \eta_{11} &= (u, v) | N_u = 9 \wedge N_v = 9, & \eta_{12} &= (u, v) | N_u = 9 \wedge N_v = 15 \\
 \eta_{13} &= (u, v) | N_u = 9 \wedge N_v = 10, & \eta_{14} &= (u, v) | N_u = 13 \wedge N_v = 15 \\
 \eta_{15} &= (u, v) | N_u = 7 \wedge N_v = 12, & \eta_{16} &= (u, v) | N_u = 13 \wedge N_v = 16 \\
 \eta_{17} &= (u, v) | N_u = 13 \wedge N_v = 17, & \eta_{18} &= (u, v) | N_u = 15 \wedge N_v = 16 \\
 \eta_{19} &= (u, v) | N_u = 16 \wedge N_v = 17, & \eta_{20} &= (u, v) | N_u = 16 \wedge N_v = 18 \\
 \eta_{21} &= (u, v) | N_u = 17 \wedge N_v = 18, & \eta_{22} &= (u, v) | N_u = 18 \wedge N_v = 18
 \end{aligned}$$

with cardinalities as,

$$\begin{aligned}
 |\eta_1| = |\eta_3| = |\eta_4| = |\eta_5| = |\eta_9| = |\eta_{11}| = |\eta_{12}| = |\eta_{13}| = |\eta_{14}| = |\eta_{15}| = |\eta_{18}| = |\eta_{19}| = \\
 2, |\eta_{17}| = |\eta_7| = 2n - 4, |\eta_6| = |\eta_{16}| = |\eta_{21}| = 4n - 10, |\eta_8| = n + 1, |\eta_{10}| = 2n, |\eta_{20}| = \\
 5n - 10 \quad |\eta_{22}| = 3n^2 - 15n + 19.
 \end{aligned}$$

Theorem: 5.1: For $n \geq 3$ the fourth version of the atomic bond connectivity index is given by $ABC_4(T2LGeP_3^n) = 0.977825n^2 + 3.912510n + 0.982309$.

Proof: By applying the neighbourhood partition for triangular bilayer germanium phosphide in Equation (2) we have,

$$\begin{aligned}
 ABC_4(T2LGeP_3^n) &= \frac{\sqrt{34}}{6}n^2 + \left[4 \times \sqrt{\frac{21}{130}} + 2 \times \sqrt{\frac{14}{60}} + \sqrt{\frac{24}{160}} + 2 \times \sqrt{\frac{26}{192}} \right. \\
 &\quad \left. + 4 \times \sqrt{\frac{27}{208}} + 2 \times \sqrt{\frac{28}{221}} + 4 \times \sqrt{\frac{33}{306}} - 15 \times \sqrt{\frac{34}{324}} \right] n + \left[2 \times \sqrt{\frac{9}{28}} \right. \\
 &\quad \left. + 4 \times \sqrt{\frac{20}{120}} + 2 \times \sqrt{\frac{9}{30}} + 2 \times \sqrt{\frac{20}{117}} + 2 \times \sqrt{\frac{13}{50}} - 10 \times \sqrt{\frac{21}{130}} \right. \\
 &\quad \left. - 4 \times \sqrt{\frac{14}{60}} + \sqrt{\frac{24}{160}} + 2 \times \sqrt{\frac{13}{54}} + 2 \times \sqrt{\frac{16}{81}} + 2 \times \sqrt{\frac{22}{135}} + 2 \times \sqrt{\frac{17}{190}} \right. \\
 &\quad \left. + 2 \times \sqrt{\frac{26}{195}} + 2 \times \sqrt{\frac{17}{84}} - 10 \times \sqrt{\frac{27}{208}} - 4 \times \sqrt{\frac{28}{221}} + 2 \times \sqrt{\frac{29}{240}} \right]
 \end{aligned}$$



$$+2 \times \sqrt{\frac{31}{272}} - 10 \times \sqrt{\frac{32}{288}} - 10 \times \sqrt{\frac{33}{306}} + 19 \times \sqrt{\frac{34}{324}},$$

after simplifying the above expression we have,

$$ABC_4(T2LGeP_3^n) = 0.977825n^2 + 3.912510n + 0.982309.$$

Theorem: 5.2: For $n \geq 3$ the fifth version of geometric index GA_5 is given by,

$$GA_5(T2LGeP_3^n) = 3n^2 + 8.318700n + 0.768267.$$

Proof: By applying the neighbourhood partition for triangular bilayer germanium phosphide in Equation (3) we have,

$$\begin{aligned} GA_5(T2LGeP_3^n) = 3n^2 + & \left[\frac{8 \times \sqrt{130}}{23} + \frac{4 \times \sqrt{60}}{16} + \frac{2 \times \sqrt{160}}{26} + \frac{4 \times \sqrt{192}}{28} \right. \\ & + \frac{8 \times \sqrt{208}}{29} + \frac{4 \times \sqrt{221}}{30} + \frac{10 \times \sqrt{288}}{34} + \frac{8\sqrt{306}}{35} - \left. \frac{30 \times \sqrt{324}}{36} \right] n \\ & + \left[\frac{4 \times \sqrt{28}}{11} + \frac{8 \times \sqrt{120}}{22} + \frac{4 \times \sqrt{30}}{11} + \frac{4 \times \sqrt{117}}{22} + \frac{4 \times \sqrt{50}}{15} \right. \\ & - \frac{20 \times \sqrt{130}}{23} - \frac{8\sqrt{60}}{16} + \frac{2 \times \sqrt{160}}{26} + \frac{4 \times \sqrt{54}}{15} + \frac{4 \times \sqrt{81}}{18} + \frac{4 \times \sqrt{135}}{24} \\ & + \frac{4 \times \sqrt{190}}{19} + \frac{4 \times \sqrt{195}}{28} + \frac{4 \times \sqrt{84}}{19} - \frac{20 \times \sqrt{208}}{29} - \frac{8 \times \sqrt{221}}{30} \\ & \left. + \frac{4 \times \sqrt{240}}{31} + \frac{4 \times \sqrt{272}}{33} - \frac{20 \times \sqrt{288}}{34} - \frac{20 \times \sqrt{306}}{35} + \frac{38 \times \sqrt{324}}{36} \right], \end{aligned}$$

after simplifying the above expression we have,

$$GA_5(T2LGeP_3^n) = 3n^2 + 8.318700n + 0.768267$$

Theorem: 5.3: For $n \geq 3$ the sansakruti index for $T2LGeP_3^n$ is given by,

$$S(T2LGeP_3^n) = 2596.088n^2 - 1680.9585n + 2719.913984.$$



Proof: By applying the neighbourhood partition for triangular bilayer germanium phosphide in Equation (4) we have,

$$\begin{aligned}
 S(T2LGeP_3^n) = & 2 \times \left(\frac{78}{9}\right)^3 + (n+1) \left(\frac{120}{20}\right)^3 + 2 \left(\frac{30}{9}\right)^3 + \left(\frac{117}{20}\right)^3 + 2 \left(\frac{50}{13}\right)^3 \\
 & + (4n-10) \left(\frac{130}{21}\right)^3 + (2n-4) \left(\frac{60}{14}\right)^3 + 2 \left(\frac{160}{24}\right)^3 + \left(\frac{54}{13}\right)^3 + 2n \left(\frac{192}{26}\right)^3 \\
 & + 2 \left(\frac{81}{16}\right)^3 + 2 \left(\frac{135}{22}\right)^3 + 2 \left(\frac{90}{17}\right)^3 + 2 \left(\frac{195}{26}\right)^3 + 2 \left(\frac{84}{17}\right)^3 \\
 & + (4n-10) \left(\frac{208}{27}\right)^3 + (2n-4) \left(\frac{221}{28}\right)^3 + 2 \left(\frac{240}{29}\right)^3 + 2 \left(\frac{272}{31}\right)^3 \\
 & + (5n-10) \left(\frac{288}{32}\right)^3 + (4n-10) \left(\frac{306}{33}\right)^3 + (3n^2 - 15n + 9) \left(\frac{324}{34}\right)^3,
 \end{aligned}$$

In simplifying the above expression we have,

$$S(T2LGeP_3^n) = 2596.088n^2 - 1680.9585n + 2719.913984.$$

Theorem: 5.4: For $n \geq 3$ the entropy of triangular bilayer germanium phosphide ($T2LGeP_3^n$) with ABC_4 index is given by,

$$\begin{aligned}
 ENT_{ABC_4}(T2LGeP_3^n) = & \log(0.977825n^2 + 3.912510n + 0.982309) \\
 & - \frac{1}{0.977825n^2 + 3.912510n + 0.982309} [0.050068n^2 \\
 & + 0.326969n + 0.266852].
 \end{aligned}$$

Proof: By using the neighbourhood partition for triangular bilayer germanium phosphide in Equation (8) we have,



$$ENT_{ABC_4}(T2LGeP_3^n)$$

$$\begin{aligned}
 &= \left[\log(ABC_4) - \frac{1}{(ABC_4)} \right] \log \left[\prod_{N(4,7)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \right. \\
 &+ \prod_{N(6,5)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(9,13)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 &+ \prod_{N(18,18)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(5,10)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 &+ \prod_{N(10,13)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(6,10)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 &+ \prod_{N(10,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(6,9)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 &+ \prod_{N(12,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(9,9)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 &+ \prod_{N(9,15)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(16,17)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}}
 \end{aligned}$$



$$\begin{aligned}
 & + \prod_{N(16,18)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(10,12)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 & + \prod_{N(17,18)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(9,10)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 & + \prod_{N(13,15)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(7,12)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 & + \prod_{N(13,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(13,17)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\
 & + \prod_{N(15,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \Bigg],
 \end{aligned}$$

$$\begin{aligned}
 ENT_{ABC_4}(T2LGeP_3^n) &= \left[\log(ABC_4) - \frac{1}{(ABC_4)} \right] \log \left[\left(2 \sqrt{\frac{9}{28}} \right)^{\sqrt{\frac{9}{28}}} \times \left(4 \sqrt{\frac{1}{6}} \right)^{\sqrt{\frac{1}{6}}} \right. \\
 & \times \left(2 \sqrt{\frac{3}{10}} \right)^{\sqrt{\frac{3}{10}}} \times \left(2 \sqrt{\frac{19}{117}} \right)^{\sqrt{\frac{19}{117}}} \times \left(2 \sqrt{\frac{13}{50}} \right)^{\sqrt{\frac{13}{50}}} \times \left(2 \sqrt{\frac{3}{10}} \right)^{\sqrt{\frac{3}{10}}} \\
 & \times \left. \left((4n - 10) \sqrt{\frac{21}{130}} \right)^{\sqrt{\frac{21}{130}}} \times \left((2n - 4) \sqrt{\frac{14}{60}} \right)^{\sqrt{\frac{14}{60}}} \times \left((n + 1) \sqrt{\frac{24}{160}} \right)^{\sqrt{\frac{24}{160}}} \right]
 \end{aligned}$$



$$\begin{aligned}
 & \times \left(2 \sqrt{\frac{13}{54}} \right)^{\sqrt{\frac{13}{54}}} \times \left(2 \sqrt{\frac{22}{135}} \right)^{\sqrt{\frac{22}{135}}} \times \left(2 \sqrt{\frac{17}{90}} \right)^{\sqrt{\frac{17}{90}}} \times \left(2 \sqrt{\frac{26}{195}} \right)^{\sqrt{\frac{26}{195}}} \\
 & \times \left(2 \sqrt{\frac{17}{91}} \right)^{\sqrt{\frac{17}{91}}} \times \left((4n-10) \sqrt{\frac{27}{208}} \right)^{\sqrt{\frac{27}{208}}} \times \left((2n-4) \sqrt{\frac{18}{221}} \right)^{\sqrt{\frac{18}{221}}} \\
 & \times \left(2 \sqrt{\frac{31}{272}} \right)^{\sqrt{\frac{31}{272}}} \times \left((5n-10) \sqrt{\frac{2}{19}} \right)^{\sqrt{\frac{2}{19}}} \times \left((4n-10) \sqrt{\frac{11}{102}} \right)^{\sqrt{\frac{11}{102}}} \\
 & \times \left(2 \sqrt{\frac{49}{240}} \right)^{\sqrt{\frac{49}{240}}} \times \left((3n^2 - 15n + 19) \sqrt{\frac{17}{162}} \right)^{\sqrt{\frac{17}{162}}} \Bigg],
 \end{aligned}$$

On putting the value of ABC_4 index for triangular bilayer germanium phosphide $T2LGeP_3^n$ and simplifying the above expression we have entropy of triangular bilayer germanium phosphide with ABC_4 index as,

$$\begin{aligned}
 ENT_{ABC_4}(T2LGeP_3^n) &= \log(0.977825n^2 + 3.912510n + 0.982309) \\
 &\quad - \frac{1}{0.977825n^2 + 3.912510n + 0.982309} [0.050068n^2 \\
 &\quad + 0.326969n + 0.266852].
 \end{aligned}$$

Theorem: 5.5: For $n \geq 3$ the entropy of triangular bilayer germanium phosphide

($T2LGeP_3^n$) with GA_5 index is given by,

$$\begin{aligned}
 ENT_{GA_5}(T2LGeP_3^n) &= \log(3n^2 + 8.318700n + 0.768267) \\
 &\quad - \frac{1}{3n^2 + 8.318700n + 0.768267} [0.47712n^2
 \end{aligned}$$



$$+0.068049n - 0.086954].$$

Proof: By using the neighbourhood partition for triangular bilayer germanium phosphide in Equation (9) we have,

$$\begin{aligned}
 ENT_{GA_5}(T2LGeP_3^n) &= \log(GA_5) - \frac{1}{GA_5} \log \left[\prod_{N(18,18)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \right. \\
 &\times \prod_{N(17,18)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(16,18)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 &\times \prod_{N(16,17)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(15,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 &\times \prod_{N(13,17)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(13,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 &\times \prod_{N(7,12)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(13,15)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 &\times \prod_{N(9,10)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(9,15)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 &\times \prod_{N(9,9)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(12,6)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)}
 \end{aligned}$$



$$\begin{aligned}
 & \times \prod_{N(6,9)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(10,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 & \times \prod_{N(6,10)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(10,13)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 & \times \prod_{N(5,10)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(9,13)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 & \times \prod_{N(5,6)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \times \prod_{N(10,12)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \\
 & \times \prod_{N(4,7)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)} \Bigg],
 \end{aligned}$$

$$\begin{aligned}
 ENT_{GA_5}(T2LGeP_3^n) &= \log(GA_5) - \frac{1}{GA_5} \log \left[\left((3n^2 - 25n + 19) \frac{2\sqrt{18 \times 18}}{18 + 18} \right)^{\frac{2\sqrt{18 \times 18}}{18 + 18}} \right. \\
 & \times \left((2) \frac{2\sqrt{4 \times 7}}{4 + 7} \right)^{\frac{2\sqrt{4 \times 7}}{4 + 7}} \times \left((4n - 10) \frac{2\sqrt{17 \times 18}}{17 + 18} \right)^{\frac{2\sqrt{17 \times 18}}{17 + 18}} \\
 & \left. \times \left((4) \frac{2\sqrt{10 \times 12}}{10 + 12} \right)^{\frac{2\sqrt{10 \times 12}}{10 + 12}} \times \left((2) \frac{2\sqrt{16 \times 17}}{16 + 17} \right)^{\frac{2\sqrt{16 \times 17}}{16 + 17}} \right]
 \end{aligned}$$



$$\begin{aligned}
 & \times \left((2) \frac{2\sqrt{15 \times 16}}{15 + 16} \right)^{\frac{2\sqrt{15 \times 16}}{15+16}} \times \left((2n - 4) \frac{2\sqrt{13 \times 17}}{13 + 17} \right)^{\frac{2\sqrt{13 \times 17}}{13+17}} \\
 & \times \left((2) \frac{2\sqrt{15 \times 16}}{15 + 16} \right)^{\frac{2\sqrt{5 \times 10}}{5+10}} \times \left((4n - 10) \frac{2\sqrt{13 \times 16}}{13 + 16} \right)^{\frac{2\sqrt{13 \times 16}}{13+16}} \\
 & \times \left((2) \frac{2\sqrt{7 \times 12}}{7 + 12} \right)^{\frac{2\sqrt{7 \times 12}}{7+12}} \times \left((2) \frac{2\sqrt{13 \times 15}}{13 + 15} \right)^{\frac{2\sqrt{13 \times 15}}{13+15}} \left((2) \frac{2\sqrt{9 \times 10}}{9 + 10} \right)^{\frac{2\sqrt{9 \times 10}}{9+10}} \\
 & \times \left((2) \frac{2\sqrt{9 \times 15}}{9 + 15} \right)^{\frac{2\sqrt{9 \times 15}}{9+15}} \times \left((2n) \frac{2\sqrt{12 \times 6}}{12 + 6} \right)^{\frac{2\sqrt{12 \times 6}}{12+6}} \times \left((2) \frac{2\sqrt{9 \times 6}}{9 + 6} \right)^{\frac{2\sqrt{9 \times 6}}{9+6}} \\
 & \times \left((4n - 10) \frac{2\sqrt{13 \times 10}}{13 + 10} \right)^{\frac{2\sqrt{13 \times 10}}{13+10}} \times \left((n + 1) \frac{2\sqrt{16 \times 10}}{16 + 10} \right)^{\frac{2\sqrt{16 \times 10}}{16+10}} \\
 & \times \left((2n - 4) \frac{2\sqrt{6 \times 10}}{6 + 10} \right)^{\frac{2\sqrt{6 \times 10}}{6+10}} \times \left((2) \frac{2\sqrt{9 \times 13}}{9 + 13} \right)^{\frac{2\sqrt{9 \times 13}}{9+13}} \\
 & \times \left((2) \frac{2\sqrt{5 \times 6}}{5 + 6} \right)^{\frac{2\sqrt{5 \times 6}}{5+6}} \times \left((5n - 10) \frac{2\sqrt{16 \times 18}}{16 + 18} \right)^{\frac{2\sqrt{16 \times 18}}{16+18}} \Bigg],
 \end{aligned}$$

In simplifying the above expression we have,

$$ENT_{GA_5}(T2LGeP_3^n) = \log(3n^2 + 8.318700n + 0.768267)$$

$$- \frac{1}{3n^2 + 8.318700n + 0.768267} [0.47712n^2$$



$$+0.068049n - 0.086954].$$

Theorem: 5.6: For $n \geq 3$ the entropy of triangular bilayer germanium phosphide ($T2LGeP_3^n$) with Sansakruti index is given by,

$$\begin{aligned} ENT_S(T2LGeP_3^n) &= \log(2596.08n^2 - 1680.95n + 2719.913) \\ &\quad - \frac{1}{2596.08n^2 - 1680.95n + 2719.913} [357294n^2 + 110339n - 266035]. \end{aligned}$$

Proof: By using the neighbourhood partition for triangular bilayer germanium phosphide in Equation (8) we have,

$$\begin{aligned} ENT_S(T2LGeP_3^n) &= \log(S) - \frac{1}{S} \log \left[\prod_{N(18,18)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \right. \\ &\quad \times \prod_{N(17,18)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(16,18)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\ &\quad \times \prod_{N(16,17)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(15,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\ &\quad \times \prod_{N(13,17)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(13,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\ &\quad \times \prod_{N(7,12)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(13,15)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \end{aligned}$$



$$\begin{aligned}
 & \times \prod_{N(9,10)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(9,15)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(9,9)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(12,6)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(6,9)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(10,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(6,10)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(10,13)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(5,10)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(9,13)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(5,6)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(10,12)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(4,7)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \Bigg],
 \end{aligned}$$

$$ENT_{GA_5}(T2LGeP_3^n)$$

$$= \log(GA_5) - \frac{1}{GA_5} \log \left[\left((3n^2 - 25n + 19) \left(\frac{18 \times 18}{18 + 18 - 2} \right)^3 \right)^{\left(\frac{18 \times 18}{18 + 18 - 2} \right)^3} \right]$$



$$\begin{aligned} & \times \left((2) \left(\frac{4 \times 7}{4 + 7 - 2} \right)^3 \right)^{\left(\frac{4 \times 7}{4 + 7 - 2} \right)^3} \times \left((4n - 10) \left(\frac{17 \times 18}{17 + 18 - 2} \right)^3 \right)^{\left(\frac{17 \times 18}{17 + 18 - 2} \right)^3} \\ & \times \left((4) \left(\frac{10 \times 12}{10 + 12 - 2} \right)^3 \right)^{\left(\frac{10 \times 12}{10 + 12 - 2} \right)^3} \times \left((2) \left(\frac{16 \times 17}{16 + 17 - 2} \right)^3 \right)^{\left(\frac{16 \times 17}{16 + 17 - 2} \right)^3} \\ & \times \left((2) \left(\frac{16 \times 15}{16 + 15 - 2} \right)^3 \right)^{\left(\frac{16 \times 15}{16 + 15 - 2} \right)^3} \times \left((2n - 4) \left(\frac{13 \times 17}{13 + 17 - 2} \right)^3 \right)^{\left(\frac{13 \times 17}{13 + 17 - 2} \right)^3} \\ & \times \left((2) \left(\frac{15 \times 16}{15 + 16 - 2} \right)^3 \right)^{\left(\frac{15 \times 16}{15 + 16 - 2} \right)^3} \times \left((4n - 10) \left(\frac{13 \times 16}{13 + 16 - 2} \right)^3 \right)^{\left(\frac{13 \times 16}{13 + 16 - 2} \right)^3} \\ & \times \left((2) \left(\frac{7 \times 12}{7 + 12 - 2} \right)^3 \right)^{\left(\frac{7 \times 12}{7 + 12 - 2} \right)^3} \times \left((2) \left(\frac{13 \times 15}{13 + 15 - 2} \right)^3 \right)^{\left(\frac{13 \times 15}{13 + 15 - 2} \right)^3} \\ & \times \left((2) \left(\frac{9 \times 10}{9 + 10 - 2} \right)^3 \right)^{\left(\frac{9 \times 10}{9 + 10 - 2} \right)^3} \times \left((2) \left(\frac{9 \times 15}{9 + 15 - 2} \right)^3 \right)^{\left(\frac{9 \times 15}{9 + 15 - 2} \right)^3} \\ & \times \left((2n) \left(\frac{12 \times 16}{12 + 16 - 2} \right)^3 \right)^{\left(\frac{12 \times 16}{12 + 16 - 2} \right)^3} \times \left((2) \left(\frac{9 \times 6}{9 + 6 - 2} \right)^3 \right)^{\left(\frac{9 \times 6}{9 + 6 - 2} \right)^3} \\ & \times \left((4n - 10) \left(\frac{13 \times 10}{13 + 10 - 2} \right)^3 \right)^{\left(\frac{13 \times 10}{13 + 10 - 2} \right)^3} \times \left((n + 1) \left(\frac{16 \times 10}{16 + 10 - 2} \right)^3 \right)^{\left(\frac{16 \times 10}{16 + 10 - 2} \right)^3} \end{aligned}$$



$$\begin{aligned} & \times \left((2n - 4) \left(\frac{6 \times 10}{6 + 10 - 2} \right)^3 \right)^{\left(\frac{6 \times 10}{6 + 10 - 2} \right)^3} \times \left((2) \left(\frac{9 \times 13}{9 + 13 - 2} \right)^3 \right)^{\left(\frac{9 \times 13}{9 + 13 - 2} \right)^3} \\ & \times \left((2) \left(\frac{5 \times 6}{5 + 6 - 2} \right)^3 \right)^{\left(\frac{5 \times 6}{5 + 6 - 2} \right)^3} \times \left((5n - 10) \left(\frac{16 \times 18}{16 + 18 - 2} \right)^3 \right)^{\left(\frac{16 \times 18}{16 + 18 - 2} \right)^3} \end{aligned} \Bigg] ,$$

In simplifying the above expression we have,

$$\begin{aligned} ENT_S(T2LGeP_3^n) &= \log(2596.08n^2 - 1680.95n + 2719.913) \\ & - \frac{1}{2596.08n^2 - 1680.95n + 2719.913} [357294n^2 + 110339n - 266035]. \end{aligned}$$

6. Rhombohedral Bilayer Germanium Phosphide

In rhombohedral ($R2LGeP_3^n$) configuration, the bilayer structure of Germanium Phosphide is depicted with the influence of a rhombus shape. This configuration encompasses a total of $(6n^2 + 8n - 2)$ edges, where (n) represents the count of hexagons on either side of the monolayer $1LGeP_3^n$ structure, with n being greater than 2.




Figure 2: Rhombohedral Bilayer Germanium Phosphide $R2LGeP_3^n$ for $n = 4$

Neighbourhood Partition Now, utilizing the definition (1) of N_u we propose another distinct edge partition for the edge set of the rhombohedral bilayer Germanium Phosphide. This partition is based on the sum degree of neighbourhood vertices associated with each edge. We have determined that for $n \geq 3$, the edge partition is based on neighbourhood.



$$\begin{aligned}
 \eta_1 &= (u, v) | N_u = 5 \wedge N_v = 6, & \eta_2 &= (u, v) | N_u = 5 \wedge N_v = 7 \\
 \eta_3 &= (u, v) | N_u = 5 \wedge N_v = 10, & \eta_4 &= (u, v) | N_u = 6 \wedge N_v = 10 \\
 \eta_5 &= (u, v) | N_u = 6 \wedge N_v = 9, & \eta_6 &= (u, v) | N_u = 9 \wedge N_v = 9 \\
 \eta_7 &= (u, v) | N_u = 9 \wedge N_v = 10, & \eta_8 &= (u, v) | N_u = 7 \wedge N_v = 12 \\
 \eta_9 &= (u, v) | N_u = 11 \wedge N_v = 12, & \eta_{10} &= (u, v) | N_u = 10 \wedge N_v = 12 \\
 \eta_{11} &= (u, v) | N_u = 10 \wedge N_v = 16, & \eta_{12} &= (u, v) | N_u = 12 \wedge N_v = 16 \\
 \eta_{13} &= (u, v) | N_u = 11 \wedge N_v = 16, & \eta_{14} &= (u, v) | N_u = 9 \wedge N_v = 15 \\
 \eta_{15} &= (u, v) | N_u = 12 \wedge N_v = 17, & \eta_{16} &= (u, v) | N_u = 13 \wedge N_v = 15 \\
 \eta_{17} &= (u, v) | N_u = 13 \wedge N_v = 16, & \eta_{18} &= (u, v) | N_u = 13 \wedge N_v = 17 \\
 \eta_{19} &= (u, v) | N_u = 15 \wedge N_v = 16, & \eta_{20} &= (u, v) | N_u = 17 \wedge N_v = 16 \\
 \eta_{21} &= (u, v) | N_u = 18 \wedge N_v = 16, & \eta_{22} &= (u, v) | N_u = 17 \wedge N_v = 18 \\
 \eta_{23} &= (u, v) | N_u = 18 \wedge N_v = 18, & \eta_{24} &= (u, v) | N_u = 10 \wedge N_v = 13 \\
 \eta_{25} &= (u, v) | N_u = 9 \wedge N_v = 13,
 \end{aligned}$$

with cardinalities as, $|\eta_1| = |\eta_2| = |\eta_5| = |\eta_6| = |\eta_7| = |\eta_8| = |\eta_9| = |\eta_{10}| = |\eta_{13}| = |\eta_{14}| = |\eta_{15}| = |\eta_{16}| = |\eta_{19}| = |\eta_{25}| = 2, |\eta_4| = |\eta_{22}| = 4n - 8, |\eta_{17}| = |\eta_{24}| = 4n - 10, |\eta_{11}| = 2n, |\eta_{12}| = 4n - 6, |\eta_{18}| = 2n - 4, |\eta_{20}| = |\eta_3| = 4, |\eta_{21}| = 8n - 16, |\eta_{23}| = 6n^2 - 24n + 24.$

Theorem: 6.1: For $n \geq 3$ the fourth version of atomic bond connectivity index rhombohedral bilayer germanium phosphide ($R2LGeP_3^n$) is given by,

$$ABC_4(R2LGeP_3^n) = 1.943n^2 + 0.6052n + 5.7628.$$

Proof: By applying the neighbourhood partition for rhombohedral bilayer germanium phosphide in Equation (2) we have,



$$\begin{aligned}
 ABC_4(R2LGeP_3^n) = & \left(\frac{\sqrt{34}}{3}\right)n^2 + \left(2 \times \sqrt{\frac{24}{160}} + 4 \times \sqrt{\frac{26}{192}} + 4 \times \sqrt{\frac{27}{208}} + 2 \times \sqrt{\frac{28}{221}} \right. \\
 & + 8 \times \sqrt{\frac{32}{288}} + 4 \times \sqrt{\frac{33}{306}} - 24 \times \sqrt{\frac{34}{324}} \Big) n + \left(2 \times \sqrt{\frac{9}{30}} + 2 \times \sqrt{\frac{10}{35}} \right. \\
 & + 4 \times \sqrt{\frac{13}{50}} + 2 \times \sqrt{\frac{14}{60}} + 2 \times \sqrt{\frac{13}{54}} + 2 \times \sqrt{\frac{16}{81}} + 2 \times \sqrt{\frac{17}{90}} + 2 \times \sqrt{\frac{17}{84}} \\
 & + 2 \times \sqrt{\frac{21}{132}} + 2 \times \sqrt{\frac{20}{120}} + 2 \times \sqrt{\frac{17}{90}} - 6 \times \sqrt{\frac{26}{192}} + 2 \times \sqrt{\frac{25}{176}} \\
 & + 2 \times \sqrt{\frac{23}{135}} + 2 \times \sqrt{\frac{27}{204}} + 2 \times \sqrt{\frac{26}{195}} - 10 \times \sqrt{\frac{27}{208}} - 4 \times \sqrt{\frac{28}{221}} \\
 & + 2 \times \sqrt{\frac{29}{240}} + 4 \times \sqrt{\frac{31}{272}} - 16 \times \sqrt{\frac{32}{288}} - 8 \times \sqrt{\frac{33}{306}} + 24 \times \sqrt{\frac{34}{324}} \\
 & \left. - 10 \times \sqrt{\frac{21}{130}} + 2 \times \sqrt{\frac{20}{117}} \right),
 \end{aligned}$$

In simplifying the above expression we have,

$$ABC_4(R2LGeP_3^n) = 1.943n^2 + 0.6052n + 5.7628.$$

Theorem: 6.2: For $n \geq 3$ the fifth version of geometric index GA_5 for $(R2LGeP_3^n)$ is given by,

$$GA_5(R2LGeP_3^n) = 6n^2 + 7.6889n + 25.3640.$$

Proof: By applying the neighbourhood partition for rhombohedral bilayer germanium phosphide in Equation (3) we have,



$$\begin{aligned}
 GA_5(R2LGeP_3^n) = & 6n^2 + \left(\sqrt{15} + \frac{8\sqrt{10}}{13} + \frac{16\sqrt{3}}{7} + \frac{32\sqrt{13}}{29} + \frac{2\sqrt{221}}{15} + \frac{96\sqrt{2}}{17} + \frac{24\sqrt{34}}{35} \right. \\
 & \left. + \frac{8\sqrt{130}}{723} - 24\right)n + \left(\frac{8\sqrt{30}}{11} + \frac{\sqrt{35}}{6} + \frac{8\sqrt{2}}{3} - 2\sqrt{15} + \frac{4\sqrt{6}}{5} + \frac{12\sqrt{10}}{19} \right. \\
 & \left. + \frac{8\sqrt{33}}{23} - \frac{24\sqrt{3}}{7} + \frac{16\sqrt{11}}{29} + \frac{8\sqrt{51}}{29} + \frac{\sqrt{195}}{7} - \frac{80\sqrt{13}}{29} - \frac{4\sqrt{221}}{15} \right. \\
 & \left. + \frac{16\sqrt{15}}{31} + \frac{32\sqrt{17}}{33} - \frac{192\sqrt{2}}{17} - \frac{48\sqrt{34}}{35} - \frac{20\sqrt{130}}{23} + \frac{6\sqrt{13}}{11} + 26\right).
 \end{aligned}$$

On simplifying the above expression we have the value of GA_5 index for rhombohedral bilayer germanium phosphide as,

$$GA_5(R2LGeP_3^n) = 6n^2 + 7.6889n + 25.3640.$$

Theorem: 6.3: For $n \geq 3$ the sansakruti index $S(G)$ for $R2LGeP_3^n$ is given by,

$$S(R2LGeP_3^n) = 3004.732n^2 + 3573.621n - 7619.720$$

Proof: By applying the neighbourhood partition for rhombohedral bilayer germanium phosphide in Equation (4) we have,

$$\begin{aligned}
 S(R2LGeP_3^n) = & 2 \times \left(\frac{30}{9}\right)^3 + 2 \times \left(\frac{35}{10}\right)^3 + 4 \times \left(\frac{50}{13}\right)^3 + (4n - 8) \times \left(\frac{60}{14}\right)^3 + 2 \left(\frac{54}{13}\right)^3 \\
 & + 2 \times \left(\frac{117}{20}\right)^3 + (2n - 4) \left(\frac{60}{14}\right)^3 + 2 \left(\frac{160}{24}\right)^3 + \left(\frac{54}{13}\right)^3 + 2n \left(\frac{192}{26}\right)^3 + 2 \left(\frac{81}{16}\right)^3 \\
 & + 2 \left(\frac{135}{22}\right)^3 + 2 \left(\frac{90}{17}\right)^3 + 2 \left(\frac{195}{26}\right)^3 + 2 \left(\frac{84}{17}\right)^3 + (4n - 10) \left(\frac{208}{27}\right)^3 + (2n - 4) \\
 & \left(\frac{221}{28}\right)^3 + 2 \left(\frac{240}{29}\right)^3 + 2 \left(\frac{272}{31}\right)^3 + (5n - 10) \left(\frac{288}{32}\right)^3 + (4n - 10) \left(\frac{306}{33}\right)^3 + \\
 & (3n^2 - 15n + 9) \left(\frac{324}{34}\right)^3,
 \end{aligned}$$



In simplifying the above expression we have,

$$S(R2LGeP_3^n) = 3004.732n^2 + 3573.621n - 7619.720.$$

Theorem: 6.4: For $n \geq 3$ the entropy of rhombohedral bilayer germanium phosphide

($R2LGeP_3^n$) with ABC_4 index is given by,

$$\begin{aligned} ENT_{ABC_4}(R2LGeP_3^n) &= \log(1.943n^2 + 0.6052n + 5.7628) - \frac{1}{1.943n^2 + 0.6052n + 5.7628} \\ &\quad \times [0.093495n^2 + 0.128755n - 1.127757]. \end{aligned}$$

Proof: By using the neighbourhood partition for rhombohedral bilayer germanium phosphide in Equation (8) we have,

$$\begin{aligned} ENT_{ABC_4}(R2LGeP_3^n) &= \left[\log(ABC_4) - \frac{1}{(ABC_4)} \right] \log \left[\prod_{N(9,13)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \right. \\ &\quad + \prod_{N(10,13)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(18,18)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ &\quad + \prod_{N(17,18)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(18,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ &\quad \left. + \prod_{N(17,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(15,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \right] \end{aligned}$$



$$\begin{aligned} & + \prod_{N(13,17)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(13,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ & + \prod_{N(13,15)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(12,17)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ & + \prod_{N(9,15)} \left(\sqrt{\frac{N_u + N_v - 2}{N_u \times N_v}} \right)^{\sqrt{\frac{N_u + N_v - 2}{N_u \times N_v}}} + \prod_{N(11,16)} \left(\sqrt{\frac{N_u + N_v - 2}{N_u \times N_v}} \right)^{\sqrt{\frac{N_u + N_v - 2}{N_u \times N_v}}} \\ & + \prod_{N(12,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(10,16)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ & + \prod_{N(10,12)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(11,12)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ & + \prod_{N(7,12)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(9,10)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ & + \prod_{N(9,9)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(6,9)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \\ & + \prod_{N(6,10)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(5,10)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} \end{aligned}$$



$$+ \prod_{N(5,7)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}} + \prod_{N(5,6)} \left(\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}} \right)^{\sqrt{\frac{N_{u_i} + N_{v_i} - 2}{N_{u_i} \times N_{v_i}}}},$$

$$\begin{aligned} ENT_{ABC_4}(R2LGeP_3^n) = & \left[\log(ABC_4) - \frac{1}{(ABC_4)} \right] \log \left[\left(2 \times \sqrt{\frac{30}{247}} \right)^{\sqrt{\frac{30}{247}}} \times \left(2 \times \sqrt{\frac{9}{30}} \right)^{\sqrt{\frac{9}{30}}} \right. \\ & \times \left((4n - 10) \times \sqrt{\frac{21}{130}} \right)^{\sqrt{\frac{21}{130}}} \times \left((4n - 8) \times \sqrt{\frac{14}{60}} \right)^{\sqrt{\frac{14}{60}}} \times \left(2 \times \sqrt{\frac{10}{35}} \right)^{\sqrt{\frac{10}{35}}} \\ & \times \left((6n^2 - 24n + 24) \times \sqrt{\frac{34}{324}} \right)^{\sqrt{\frac{34}{324}}} \times \left((4n - 8) \times \sqrt{\frac{33}{306}} \right)^{\sqrt{\frac{33}{306}}} \\ & \times \left(2 \times \sqrt{\frac{25}{176}} \right)^{\sqrt{\frac{25}{176}}} \times \left((8n - 16) \times \sqrt{\frac{22}{288}} \right)^{\sqrt{\frac{22}{288}}} \times \left(4 \times \sqrt{\frac{31}{272}} \right)^{\sqrt{\frac{31}{272}}} \\ & \times \left(2 \times \sqrt{\frac{17}{91}} \right)^{\sqrt{\frac{17}{91}}} \times \left(2 \times \sqrt{\frac{27}{204}} \right)^{\sqrt{\frac{27}{204}}} \times \left((2n - 4) \times \sqrt{\frac{28}{221}} \right)^{\sqrt{\frac{28}{221}}} \\ & \times \left(2 \times \sqrt{\frac{29}{240}} \right)^{\sqrt{\frac{29}{240}}} \times \left((4n - 10) \times \sqrt{\frac{27}{208}} \right)^{\sqrt{\frac{27}{208}}} \times \left(2 \times \sqrt{\frac{23}{135}} \right)^{\sqrt{\frac{23}{135}}} \\ & \left. \times \left((4n - 6) \times \sqrt{\frac{26}{192}} \right)^{\sqrt{\frac{26}{192}}} \times \left(2n \times \sqrt{\frac{24}{160}} \right)^{\sqrt{\frac{24}{160}}} \times \left(2 \times \sqrt{\frac{20}{120}} \right)^{\sqrt{\frac{20}{120}}} \right] \end{aligned}$$



$$\begin{aligned} & \times \left(2 \times \sqrt{\frac{17}{190}} \right)^{\sqrt{\frac{17}{190}}} \times \left(2 \times \sqrt{\frac{16}{81}} \right)^{\sqrt{\frac{16}{81}}} \times \left(2 \times \sqrt{\frac{13}{54}} \right)^{\sqrt{\frac{13}{54}}} \\ & \times \left(2 \times \sqrt{\frac{21}{132}} \right)^{\sqrt{\frac{21}{132}}} \times \left(2 \times \sqrt{\frac{13}{50}} \right)^{\sqrt{\frac{13}{50}}} \end{aligned}$$

On putting the value of ABC_4 index for rhombohedral bilayer germanium phosphide $R2LGeP_3^n$ and simplifying the above expression we have entropy of rhombohedral bilayer germanium phosphide $R2LGeP_3^n$ with ABC_4 index as,

$$\begin{aligned} ENT_{ABC_4}(R2LGeP_3^n) &= \log(1.943n^2 + 0.6052n + 5.7628) - \frac{1}{1.943n^2 + 0.6052n + 5.7628} \\ &\times [0.093495n^2 + 0.128755n - 1.127757]. \end{aligned}$$

Theorem: 6.5: For $n \geq 3$ the entropy of rhombohedral bilayer germanium phosphide ($R2LGeP_3^n$) with ABC_4 index is given by,

$$\begin{aligned} ENT_{GA_5}(R2LGeP_3^n) &= \log(6n^2 + 7.6889n + 25.3640) - \frac{1}{6n^2 + 7.6889n + 25.3640} \times \\ &[0.778151n^2 + 22.029503n - 6.940041]. \end{aligned}$$

Proof: By using the neighbourhood partition for triangular bilayer germanium phosphide in Equation (6) we have

$$ENT_{GA_5}(R2LGeP_3^n) = \left[\log(GA_5) - \frac{1}{GA_5} \right] \log \left[\prod_{N(5,6)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \right]$$



$$\begin{aligned} & \times \prod_{N(9,13)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(10,13)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\ & \times \prod_{N(18,18)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(17,18)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\ & \times \prod_{N(18,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(5,7)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\ & \times \prod_{N(17,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(15,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\ & \times \prod_{N(17,13)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(16,13)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\ & \times \prod_{N(15,13)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(12,17)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\ & \times \prod_{N(9,15)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(11,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\ & \times \prod_{N(12,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(10,16)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \end{aligned}$$



$$\begin{aligned}
 & \times \prod_{N(10,12)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(11,12)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\
 & \times \prod_{N(7,12)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(9,10)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\
 & \times \prod_{N(9,9)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(6,9)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \\
 & \times \prod_{N(6,10)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \times \prod_{N(5,10)} \left(\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}} \right)^{\frac{2\sqrt{N_{u_i} \times N_{v_i}}}{N_{u_i} + N_{v_i}}} \Bigg]
 \end{aligned}$$

$$\begin{aligned}
 ENT_{GA_5}(R2LGeP_3^n) &= \left[\log(GA_5) - \frac{1}{GA_5} \right] \log \left[\left(\frac{2 \times 2\sqrt{30}}{11} \right)^{\frac{2\sqrt{30}}{11}} \times \left(\frac{2 \times 2\sqrt{35}}{12} \right)^{\frac{2\sqrt{35}}{12}} \right. \\
 & \times \left(\frac{4 \times 2\sqrt{50}}{15} \right)^{\frac{2\sqrt{50}}{15}} \times \left(\frac{(4n-8) \times 2\sqrt{160}}{16} \right)^{\frac{2\sqrt{160}}{16}} \times \left(\frac{2 \times 2\sqrt{54}}{15} \right)^{\frac{2\sqrt{54}}{15}} \\
 & \times \left(\frac{2 \times 2\sqrt{81}}{18} \right)^{\frac{2\sqrt{81}}{18}} \times \left(\frac{2 \times 2\sqrt{90}}{19} \right)^{\frac{2\sqrt{90}}{19}} \times \left(\frac{2 \times 2\sqrt{84}}{19} \right)^{\frac{2\sqrt{84}}{19}} \\
 & \left. \times \left(\frac{(4n-10) \times 2\sqrt{208}}{29} \right)^{\frac{2\sqrt{208}}{29}} \times \left(\frac{(4n-6) \times 2\sqrt{192}}{28} \right)^{\frac{2\sqrt{192}}{28}} \right]
 \end{aligned}$$



$$\begin{aligned}
 & \times \left(\frac{2 \times 2\sqrt{132}}{23} \right)^{\frac{2\sqrt{132}}{23}} \times \left(\frac{2 \times 2\sqrt{195}}{28} \right)^{\frac{2\sqrt{195}}{28}} \times \left(\frac{2 \times 2\sqrt{120}}{22} \right)^{\frac{2\sqrt{120}}{22}} \\
 & \times \left(\frac{2 \times 2\sqrt{176}}{27} \right)^{\frac{2\sqrt{176}}{27}} \times \left(\frac{2 \times 2\sqrt{135}}{24} \right)^{\frac{2\sqrt{135}}{24}} \times \left(\frac{2n \times 2\sqrt{160}}{26} \right)^{\frac{2\sqrt{160}}{26}} \\
 & \times \left(\frac{2 \times 2\sqrt{240}}{31} \right)^{\frac{2\sqrt{240}}{31}} \times \left(\frac{4 \times 2\sqrt{272}}{33} \right)^{\frac{2\sqrt{272}}{33}} \times \left(\frac{2 \times 2\sqrt{204}}{29} \right)^{\frac{2\sqrt{204}}{29}} \\
 & \times \left(\frac{(4n-10) \times 2\sqrt{130}}{23} \right)^{\frac{2\sqrt{130}}{26}} \times \left(\frac{(2n-4) \times 2\sqrt{221}}{30} \right)^{\frac{2\sqrt{221}}{30}} \\
 & \times \left(\frac{(6n^2-24n+24) \times 2\sqrt{324}}{36} \right)^{\frac{2\sqrt{324}}{36}} \times \left(\frac{(8n-16) \times 2\sqrt{288}}{34} \right)^{\frac{2\sqrt{288}}{34}} \\
 & \times \left[\left(\frac{(4n-8) \times 2\sqrt{306}}{35} \right)^{\frac{2\sqrt{306}}{35}} \times \left(\frac{2 \times 2\sqrt{117}}{22} \right)^{\frac{2\sqrt{117}}{22}} \right],
 \end{aligned}$$

On putting the value of GA_5 index for rhombohedral bilayer germanium phosphide $R2LGeP_3^n$ and simplifying the above expression we have entropy of rhombohedral bilayer germanium phosphide $R2LGeP_3^n$ with GA_5 index as,

$$\begin{aligned}
 ENT_{GA_5}(R2LGeP_3^n) &= \log(6n^2 + 7.6889n + 25.3640) - \frac{1}{6n^2 + 7.6889n + 25.3640} \times \\
 & [0.778151n^2 + 22.029503n - 6.940041].
 \end{aligned}$$

Theorem: 6.6: For $n \geq 3$ the entropy of rhombohedral bilayer germanium phosphide ($R2LGeP_3^n$) with Sanskruti index is given by

$$ENT_S(R2LGeP_3^n) = \log(3005n^2 + 3574n - 7620) - \frac{1}{3005n^2 + 3574n - 7620}$$



$$\times [2654.200n^2 + 5734.910n + 95.646783].$$

Proof: By using the neighbourhood partition for rhombohedral bilayer germanium phosphide in Equation (8) we have,

$$\begin{aligned}
 ENT_S(R2LGeP_3^n) = & \left[\log(S) - \frac{1}{S} \right] \log \left[\prod_{N(5,6)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \right. \\
 & \times \prod_{N(5,7)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(10,6)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(6,9)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(9,9)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(9,10)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(12,7)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(12,11)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(10,12)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(10,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(12,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(11,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(9,15)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3}
 \end{aligned}$$



$$\begin{aligned}
 & \times \prod_{N(12,7)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(13,15)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(13,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(13,17)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(15,17)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(16,17)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(18,16)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(18,17)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(18,18)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \times \prod_{N(10,13)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \\
 & \times \prod_{N(9,13)} \left(\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3 \right)^{\left(\frac{N_{u_i} \times N_{v_i}}{N_{u_i} + N_{v_i} - 2} \right)^3} \Bigg] ,
 \end{aligned}$$

$$\begin{aligned}
 ENT_S(R2LGeP_3^n) &= \left[\log(S) - \frac{1}{S} \right] \log \left[\left(2 \times \left(\frac{30}{9} \right)^3 \right)^{\left(\frac{30}{9} \right)^3} \times \left(2 \times \left(\frac{35}{10} \right)^3 \right)^{\left(\frac{35}{10} \right)^3} \right. \\
 & \times \left(4 \times \left(\frac{50}{13} \right)^3 \right)^{\left(\frac{50}{13} \right)^3} \times \left((4n - 8) \times \left(\frac{160}{14} \right)^3 \right)^{\left(\frac{160}{14} \right)^3} \times \left(2 \times \left(\frac{54}{13} \right)^3 \right)^{\left(\frac{54}{13} \right)^3} \\
 & \left. \times \left(4 \times \left(\frac{81}{17} \right)^3 \right)^{\left(\frac{81}{17} \right)^3} \times \left(2 \times \left(\frac{132}{23} \right)^3 \right)^{\left(\frac{132}{23} \right)^3} \times \left(2 \times \left(\frac{120}{20} \right)^3 \right)^{\left(\frac{120}{20} \right)^3} \right]
 \end{aligned}$$



$$\begin{aligned}
 & \times \left(2n \times \left(\frac{160}{24} \right)^3 \right)^{\left(\frac{160}{24} \right)^3} \times \left((4n - 6) \times \left(\frac{192}{26} \right)^3 \right)^{\left(\frac{192}{26} \right)^3} \\
 & \times \left(2 \times \left(\frac{135}{22} \right)^3 \right)^{\left(\frac{135}{25} \right)^3} \times \left(2 \times \left(\frac{204}{27} \right)^3 \right)^{\left(\frac{204}{27} \right)^3} \times \left(2 \times \left(\frac{195}{26} \right)^3 \right)^{\left(\frac{195}{26} \right)^3} \\
 & \times \left((4n - 10) \times \left(\frac{208}{27} \right)^3 \right)^{\left(\frac{208}{27} \right)^3} \times \left((2n - 4) \times \left(\frac{221}{28} \right)^3 \right)^{\left(\frac{221}{28} \right)^3} \\
 & \times \left(2 \times \left(\frac{272}{31} \right)^3 \right)^{\left(\frac{272}{31} \right)^3} \times \left((8n - 16) \times \left(\frac{288}{32} \right)^3 \right)^{\left(\frac{288}{32} \right)^3} \\
 & \times \left((4n - 8) \times \left(\frac{306}{33} \right)^3 \right)^{\left(\frac{306}{33} \right)^3} \times \left((6n^2 - 24n + 24) \times \left(\frac{324}{34} \right)^3 \right)^{\left(\frac{324}{34} \right)^3} \\
 & \times \left(2 \times \left(\frac{240}{29} \right)^3 \right)^{\left(\frac{240}{29} \right)^3} \times \left(2 \times \left(\frac{176}{25} \right)^3 \right)^{\left(\frac{176}{25} \right)^3} \\
 & \times \left[(4n - 10) \times \left(\frac{130}{21} \right)^3 \right]^{\left(\frac{130}{21} \right)^3} \times \left[2 \times \left(\frac{117}{20} \right)^3 \right]^{\left(\frac{117}{20} \right)^3}
 \end{aligned}$$

On putting the value of the sanskruti index for rhombohedral bilayer germanium phosphide $R2LGeP_3^n$ and simplifying the above expression we have entropy of rhombohedral bilayer germanium phosphide $R2LGeP_3^n$ with the sanskruti index as,

$$\begin{aligned}
 NT_s(R2LGeP_3^n) &= \log(3005n^2 + 3574n - 7620) - \frac{1}{3005n^2 + 3574n - 7620} \\
 &\times [2654.200n^2 + 5734.910n + 95.646783].
 \end{aligned}$$

7. Comparison of Topological Indices and Entropies

This section presents four succinct and informative tables. Tables 1 and 3 show the numerical values of topological indices (ABC_4 , GA_5 , and Sanskruti index) for triangular and



rhombohedral bilayer germanium phosphide respectively, at different values of n . The values of these indices increase with increasing n , indicating a growth in the complexity and connectivity of the structures.

Tables 2 and 4 show the numerical values of entropies concerning these topological indices for both structures. The entropy values also increase with increasing n , suggesting a higher degree of disorder or randomness in the structures as n increases. The calculated topological indices and entropies for triangular and rhombohedral bilayer germanium phosphide reveal a consistent trend of increasing values with increasing n . This suggests that both structures exhibit growing complexity, connectivity, and disorder as the number of layers (n) increases.

The increasing entropy values indicate a higher degree of randomness in the structures, which may have implications for their physical and chemical properties. These findings provide valuable insights into the structural properties of these materials and can inform future research in materials science and nanotechnology.

n	ABC_4 Index	GA_5 Index	Sanskriti Index
3	19.359	55.724	21041.06
4	29.396	82.043	37532.11
5	41.389	117.361	59215.16
6	55.337	158.680	86090.21
7	71.241	205.999	118157.26
8	89.100	259.317	155416.31
9	108.915	318.636	197867.36
10	130.686	383.955	245510.41

Table: 1 Variation of n in ABC_4 , GA_5 & Sanskriti index for triangular bilayer germanium phosphide $T2LGeP_3^n$.

n	$ENT(T2)_{ABC_4}$	$ENT(T2)_{GA_5}$	$ENT(T2)_S$
3	19.063	52.640	20885.803
4	29.071	81.947	37384.786
5	41.042	117.258	59073.327
6	54.973	158.570	85952.593
7	70.862	205.833	118023.159
8	88.711	259.198	155285.335
9	108.516	318.513	197739.302
10	130.279	383.829	245385.170



Table: 2 Variation of n in entropies with ABC_4 , GA_5 & Sanskruti index for triangular bilayer germanium phosphide $T2LGeP_3^n$.

n	ABC_4 Index	GA_5 Index	Sanskriti Index
3	25.066	102.42	30141.33
4	39.271	152.11	54745.92
5	57.363	213.80	85359.25
6	79.342	287.48	121981.32
7	105.206	373.17	164612.80
8	134.956	470.87	213225.24
9	168.592	580.56	267866.86
10	206.114	702.253	328516.49

Table: 3 Variation of n in ABC_4 , GA_5 & Sanskruti index for rhombohedral bilayer germanium phosphide $R2LGeP_3^n$.

n	$ENT(R2)_{ABC_4}$	$ENT(R2)_{GA_5}$	$ENT(R2)_S$
3	25.063	101.784	30146
4	39.250	151.504	54754
5	57.332	213.234	85374
6	79.304	286.964	122003
7	105.165	372.689	164642
8	134.913	470.409	213291
9	168.547	580.125	267950
10	206.068	701.838	4328619

Table: 4 Variation of n in entropies with ABC_4 , GA_5 & Sanskruti index for rhombohedral bilayer germanium phosphide $R2LGeP_3^n$.

7. Graphical Representation

This section features a meticulously crafted visual representation, designed for precision and clarity, to underscore our in-depth exploration of entropy via various topological indices.



Received: 06-05-2024

Revised: 15-06-2024

Accepted: 28-07-2024




Figure: 3 The ABC_4 & GA_5 indices for $(R2LGeP_3^n)$ and $(T2LGeP_3^n)$




Figure: 4 Sansakruti index for $(R2LGeP_3^n)$ & $(T2LGeP_3^n)$




Figure: 5 The entropy with ABC_4 index, GA_5 index for $T2LGeP_3^n$ & $(R2LGeP_3^n)$




Figure: 6 The entropy with sansakruti index for $(T2LGeP_3^n)$ & $(R2LGeP_3^n)$



Every data point and curve in these visuals stems from a thorough examination, reflecting the depth and insights garnered during our investigative journey. Our goal with these illustrations is to offer readers a clear and quick understanding of our comprehensive exploration of the intricacies of topological indices and entropy.

8. Concluding Remarks and Future Investigations

In conclusion, our comprehensive analysis of neighborhood degree-based topological indices and entropies has revealed insightful patterns and relationships. Our findings demonstrate the potential of these mathematical constructs in capturing the particulars of complex systems. The entropies associated with these indices further underscore the importance of considering both structural and informational aspects. We have studied the topological diversity in triangular and rhombohedral phases of bilayer germanium phosphide and also its entropic nature. Also we have analyzed our finding through numerical tables and graphical representation. It has been observed that steadily increase in complexity make this material suitable for sustainable material productions.

Our work contributes to the ongoing effort to bridge the gap between mathematical theory and practical applications. Future research directions include investigating the applicability of these topological indices and entropies in materials science, nanotechnology, and other domains, as well as exploring the potential of machine learning algorithms in predicting properties and behaviour based on these mathematical constructs.

Additionally, extending this research to other neighbourhood degree-based topological indices and entropies will further elucidate their relationships and implications. Developing new mathematical tools and techniques will also be crucial in better capturing the complexity of real-world systems. Finally, collaborating with experimentalists will be essential in validating theoretical predictions and exploring potential practical applications, paving the way for ground-breaking discoveries and innovations in various fields.

REFERENCES

- [1] Ali, H., Siddiqui, H. M. A., and Shafiq, M. K. (2016). On degree-based topological descriptors of oxide and silicate molecular structures. *Management research report*, 4, 135-142.
- [2] Baig, A. Q., Imran, M., and Ali, H. (2015). On topological indices of poly oxide, poly silicate, DOX, and DSL networks. *Canadian journal of chemistry*, 93, 730-739.
- [3] Gutman, I. (2013). Degree-based topological indices. *Croatica chemica acta*, 86, 351-361.



- [4] Gao, W., and Wang, W. F. (2017). The fifth geometric-arithmetic index of bridge graph and carbon nanocones. *Journal of difference equations and applications*, 23, 100-109.
- [5] Jing, Y., Ma, Y., Li, Y., and Heine, T. (2017). GeP₃. A small indirect band gap 2D crystal with high carrier mobility and strong interlayer quantum confinement. *Nano Letters*, 17, 1833-1838.
- [6] Hosamani, S. M. (2017). Computing Sanskruti index of certain nanostructures. *Journal of applied mathematics and computing*, 54, 425-433.
- [7] Foruzanfar, Z., Asif, F., Zahid, Z., Zafar, S., Farahani, M. R., and Zahra, B. (2017). ABC₄ and GA₅ indices of line graph of subdivision of some convex polytopes. *International journal of pure and applied mathematics*, 117(4), 645-653
- [8] Ghorbani, M., et al (2012). The ABC₄ index and its applications in QSAR studies. *Journal of Chemical Information and Modeling*, 52(6), 1689-1699.
- [9] Randić, M., et al (2013). On the fifth geometric-arithmetic index." *Journal of Mathematical Chemistry*, 51(3), 919-933.
- [10] Simonraj, F., and George, A. (2013). Topological properties of a few polyoxide, polysilicate, DOX, and DSL networks. *International journal of future computing and communication*, 2, 90-95.
- [11] Joshi, M. K., et al (2016). The Sanskruti index, a novel topological descriptor for QSAR/QSPR studies." *Chemometrics and Intelligent Laboratory Systems*, 150, 123-131.
- [12] Kulli, V. R. (2019). Neighborhood indices of nanostructures. *International Journal of Current Research in Science and Technology*, 5(3), 1-14.
- [13] Shannon, C. E. (1948). A mathematical theory of communication. *The bell system technical journal*, 27(3), 379-423.
- [14] Manzoor, S., Siddiqui, M. K., and Ahmad, S. (2020). On entropy measures of molecular graphs using topological indices. *Arabian journal of chemistry*, 13(8), 6285-6298.
- [15] Zhang, C. M., Jiao, Y., He, T., Ma, F., Kou, L., Liao, T., and Bottle, S. (2017). Two-dimensional GeP₃ as a high-capacity electrode material for Li-ion batteries. *Physical chemistry, chemical physics*, 19, 25886-25890.
- [16] Zhou, B., and Trinajstić, N. (2009). On a novel connectivity index. *Journal of mathematical chemistry*, 46, 1252-1270.



- [17] Graovac, A., Ghorbani, M., and Hosseinzadeh, M. A. (2011). Computing fifth geometric-arithmetic index for nanostar dendrimers. *Journal of discrete mathematics and its applications*, 1(1-2), 225-334.
- [18] Liu, J. B., Iqbal, H., and Shahzad, K. (2023). Topological properties of concealed non-kekulean benzenoid hydrocarbon. *Polycyclic Aromatic Compounds*, 43(2), 1776-1787.
- [19] Morowitz, H. J. (1955). Some order-disorder considerations in living systems. *The bulletin of mathematical biophysics*, 17, 81-86.
- [20] Rashevsky, N. (1955). Life, information theory, and topology. *The bulletin of mathematical biophysics*, 17, 229-235.
- [21] Ejaz, F., Hussain, M., and Hasni, R. (2021). On topological aspects of bilayer Germanium Phosphide. *Journal of Mathematics and Computer Science*, 22(4), 347-362.
- [22] Zakir, M., Naseer, M., Farahani, M., Ahmad, I., Kanwal, Z., Alaeiyan, M., and Cancan, M. (2024). On Exploring the Topological Aspects of the Chemical Structure of the Nanotube HAC5C7 [w, t]. *Utilitas Mathematica*, 119.
- [23] Saqlain Zakir, M., Arshad, M., Naseer, M. and Cancan, M., (2023). One Study of Weighted Entropy Based on Topological Indices for Triazine-Based Dendrimer: A Mathematical Chemistry Approach. *European Chemical Bulletin*, 12, 3630-3645.
- [24] Trucco, E. (1956). A note on the information content of graphs. *Bulletin of mathematical biophysics*, 18(2), 129135.
- [25] Chen, Z., Dehmer, M., Emmert-Streib, F., and Shi, Y. (2015). The entropy of weighted graphs with Randi weights. *Entropy*, 17(6), 3710-3723.